

APC Rabbit anti-Human CD271/NGFR mAb

Catalog No.: A28359

Basic Information

Observed MW

Calculated MW

35kDa/45kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC57800

Conjugate

APC. Ex:650nm. Em:660nm.

Background

Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain.

Recommended Dilutions

FC 5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

4804

Swiss Prot

P08138

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 K-562 cells (negative control, left) and SK-N-MC cells (right) were surface-stained with APC Rabbit anti-Human CD271/NGFR mAb (A28359, 5 μ l/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 SK-N-MC cells were surface-stained with APC Rabbit IgG isotype control (A24173, 5 μ l/Test, left) or APC Rabbit anti-Human CD271/NGFR mAb (A28359, 5 μ l/Test, right).