

ZO-1 Rabbit mAb

Catalog No.: A28491 **Recombinant**

Basic Information

Observed MW

250 kDa

Calculated MW

195 kDa

Category

Primary antibody

Applications

WB,IF/ICC,IF-F,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC77690

Background

Enables connexin binding activity and protein domain specific binding activity. Involved in several processes, including ameloblast differentiation; negative regulation of stress fiber assembly; and positive regulation of sprouting angiogenesis. Acts upstream of or within blastocyst formation and sensory perception of sound. Located in several cellular components, including apicolateral plasma membrane; bicellular tight junction; and intercellular canaliculus. Part of protein-containing complex. Is expressed in several structures, including alimentary system; central nervous system; early conceptus; genitourinary system; and sensory organ. Orthologous to human TJP1 (tight junction protein 1).

Recommended Dilutions

WB 1:2000 - 1:5000**IF/ICC** 1:200 - 1:800**IF-F** 1:200 - 1:800**IHC-P** 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID

21872

Swiss Prot

P39447

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ZO1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Confocal imaging of mIMCD-3 cells using ZO-1 Rabbit mAb (A28491, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of MCF7 cells using ZO-1 Rabbit mAb (A28491, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of frozen sections Mouse colon using [KO Validated] ZO-1 Rabbit mAb (A28491, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of frozen sections Mouse kidney using [KO Validated] ZO-1 Rabbit mAb (A28491, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of frozen sections Rat kidney using ZO-1 Rabbit mAb (A28491, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human pancreas tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure antigen




Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure



Immunohistochemistry analysis of paraffin-embedded Mouse intestine tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval


Validation Data

retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat pancreas tissue using ZO-1 Rabbit mAb (A28491) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.