

Anti-Monkey CD14 mAb

Catalog No.: A28754

Basic Information

Observed MW

Calculated MW

40 kDa

Category

Primary antibody

Applications

IHC-P, mIHC

Cross-Reactivity

Cynomolgus monkey

CloneNo number

ARC5278-01

Background

The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide, and to viruses. This gene has been identified as a target candidate in the treatment of SARS-CoV-2-infected patients to potentially lessen or inhibit a severe inflammatory response. Alternative splicing results in multiple transcript variants encoding the same protein.

Recommended Dilutions

IHC-P 1:200 - 1:1000

mIHC 1:100 - 1:400

Immunogen Information

Gene ID

102129342

Swiss Prot

B3Y6B7

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD14

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Cynomolgus monkey spleen tissue using Anti-Monkey CD14 mAb (A28754) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



The multiplex IHC analysis on paraffin-embedded Cynomolgus monkey spleen tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : Anti-Monkey CD4 PolymAb® (A28749PM, 1:400) with TSA-CFP440 (Light Brown), and Anti-Monkey CD8A Rabbit PolymAb® (A28747PM, 1:1000) with TSA-CFP490 (Green), and Anti-Monkey CD3 mAb (A28764, 1:1000) with TSA-CFP555 (Red), and Anti-Monkey CD20 mAb (A28759, 1:1000) with TSA-CFP645 (Yellow), and Anti-Monkey CD163 PolymAb® (A28748PM, 1:20000) with TSA-CFP440 (Magenta), and Anti-Monkey CD19 mAb (A28753, 1:400) with TSA-CFP490 (Orange), and Anti-Monkey CD14 mAb (A28754, 1:200) with TSA-CFP555 (Magenta), and Anti-Monkey CD16 mAb (A28756, 1:5000) with TSA-CFP649 (Cyan). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M citrate buffer at pH 6.0. The analysis was completed using a 40x objective lens.