

Anti-Monkey Olig2 mAb

Catalog No.: A28765

Basic Information

Observed MW

Calculated MW

32 kDa

Category

Primary antibody

Applications

IHC-P

Cross-Reactivity

Cynomolgus monkey

CloneNo number

ARC5382-01

Background

This gene encodes a basic helix-loop-helix transcription factor which is expressed in oligodendroglial tumors of the brain. The protein is an essential regulator of ventral neuroectodermal progenitor cell fate. The gene is involved in a chromosomal translocation t(14;21)(q11.2;q22) associated with T-cell acute lymphoblastic leukemia. Its chromosomal location is within a region of chromosome 21 which has been suggested to play a role in learning deficits associated with Down syndrome.

Recommended Dilutions

IHC-P 1:200 - 1:800

Immunogen Information

Gene ID

10215

Swiss Prot

Q13516

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

BHLHB1; OLIGO2; RACK17; PRKCBP2; bHLHe19

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide, 0.05% BSA, 50% glycerol, pH7.3

Validation Data



Immunohistochemistry analysis of paraffin-embedded Cynomolgus monkey brain tissue using Anti-Monkey Olig2 mAb (A28765) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



The multiplex IHC analysis on paraffin-embedded Cynomolgus monkey brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : Anti-Monkey NeuN mAb (A28655, 1:4000) with TSA-CFP440 (Magenta), and Anti-Monkey Olig2 mAb (A28765, 1:400) with TSA-CFP515 (Green), and Anti-Monkey GFAP mAb (A28763, 1:20000) with TSA-CFP555 (Red), and Anti-Monkey AIF1/IBA1 mAb (A28653, 1:10000) with TSA-CFP645 (Yellow). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M citrate buffer at pH 6.0. The analysis was completed using a 40x objective lens.