

PE Rabbit anti-Human CD178/Fas Ligand mAb

Catalog No.: A29377

Basic Information

Observed MW**Calculated MW****Category**

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC67683

Conjugate

PE. Ex:565nm. Em:574nm.

Background

This gene is a member of the tumor necrosis factor superfamily. The primary function of the encoded transmembrane protein is the induction of apoptosis triggered by binding to FAS. The FAS/FASLG signaling pathway is essential for immune system regulation, including activation-induced cell death (AICD) of T cells and cytotoxic T lymphocyte induced cell death. It has also been implicated in the progression of several cancers. Defects in this gene may be related to some cases of systemic lupus erythematosus (SLE). Alternatively spliced transcript variants have been described.

Recommended Dilutions

FC 5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

356

Swiss Prot

P48023

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

APTL; FASL; CD178; CD95L; ALPS1B; CD95-L; TNFSF6; TNLG1A; APT1LG1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 Human PBMC (untreated, left) and Human PBMC (treated with 50 ng/mL PMA and 1 µg/mL Ionomycin for 6 hours, right) were surface-stained with ABflo® 450 Rabbit anti-Human/Monkey CD3 mAb (A27177, 5 µl/Test), ABflo® 647 Rabbit anti-Human/Monkey CD69 mAb (A25892, 5 µl/Test) and PE Rabbit anti-Human CD178/Fas Ligand mAb (A29377, 5 µl/Test). Cells in the CD3⁺ gate were used for analysis.