

VEGF Receptor 2 Rabbit mAb

Catalog No.: A29388 **Recombinant**

Basic Information

Observed MW

210 kDa/230 kDa

Calculated MW

75 kDa/79 kDa/152 kDa

Category

Primary antibody

Applications

WB, IF-F, IHC-P, ELISA

Cross-Reactivity

Mouse, Rat

CloneNo number

ARC3893

Background

Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin α V β 3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.

Recommended Dilutions

WB 1:10000 - 1:40000**IF-F** 1:1000 - 1:6000**IHC-P** 1:1000 - 1:4000

ELISA Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ($\geq 1:10000$) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn

Immunogen Information

Gene ID

3791

Swiss Prot

P35968

Immunogen

This information is considered to be commercially sensitive.

Synonyms

FLK1; CD309; VEGFR; VEGFR2

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification


Storage

Store at -20°C



www.abclonal.com.cn

Validation Data



Western blot analysis of various lysates using VEGF Receptor 2 Rabbit mAb (A29388) at 1:20000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30 s.




Confocal imaging of frozen sections of Mouse pancreas tissue using VEGF Receptor 2 Rabbit mAb (A29388, dilution 1:5000) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Rat pancreas tissue using VEGF Receptor 2 Rabbit mAb (A29388, dilution 1:5000) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Immunohistochemistry analysis of paraffin-embedded Mouse heart tissue using VEGF Receptor 2 Rabbit mAb (A29388) at a dilution of 1:1400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using VEGF Receptor 2 Rabbit mAb (A29388) at a dilution of 1:1400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.