

# Histone H2A Rabbit mAb

Catalog No.: A29391 **Recombinant**

## Basic Information

**Observed MW**

14 kDa

**Calculated MW**

14 kDa

**Category**

Primary antibody

**Applications**

WB, IF/ICC, ChIP, ELISA

**Cross-Reactivity**

Human, Mouse, Rat, Monkey

**CloneNo number**

ARC4000

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

## Recommended Dilutions

**WB** 1:1000 -  
1:10000(Human&Monkey)  
1:10000 -  
1:50000(Mouse&Rat)

**IF/ICC** 1:300 - 1:600

**ChIP** 2 µg antibody for 10 µg - 15  
µg of Chromatin

**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on your  
specific assay requirements.  
For high-ratio antibody  
dilutions ( $\geq 1:10000$ ) a  
sequential dilution method  
is strongly recommended to  
ensure measurement  
accuracy.

## Immunogen Information

**Gene ID**

3012/8329

**Swiss Prot**

P04908/P0C0S8

**Immunogen**

This information is considered to be commercially sensitive.

**Synonyms**

H2A.1; H2A.2; H2A/a; H2AC4; H2AFA; HIST1H2AE; Histone H2A

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**

Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Contact

---




☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)




🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

---

Validation Data



Chromatin immunoprecipitation was performed with 10 µg of cross-linked chromatin from HeLa cells, using 2 µg of Histone H2A Rabbit mAb (A29391) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.



Validation Data

---

|                                                                                                                                                                                                  |                                                                                                                                                                                                  |                                                                                                                                                                                                  |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x. | (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x. | (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x. |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|