

[KO Validated] SMAD2/SMAD3 Rabbit mAb

Catalog No.: A29434 **KO** **Validated** **Recombinant**

Basic Information

Observed MW

52 kDa(SMAD3)/60 kDa(SMAD2)

Calculated MW

25-52 kDa

Category

Primary antibody

Applications

WB,IP,IF/ICC,ChIP,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC3701

Background

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. The SMAD family of proteins are a group of intracellular signal transducer proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. The SMAD3 protein functions in the transforming growth factor-beta signaling pathway, and transmits signals from the cell surface to the nucleus, regulating gene activity and cell proliferation. This protein forms a complex with other SMAD proteins and binds DNA, functioning both as a transcription factor and tumor suppressor. Mutations in this gene are associated with aneurysms-osteoarthritis syndrome and Loews-Dietz Syndrome 3.

Recommended Dilutions

WB 1:7000 - 1:40000

IP 0.5 µg - 4 µg antibody for
200 µg - 400 µg extracts of
whole cells

IF/ICC 1:200 - 1:1400

ChIP 1 µg antibody for 10 µg - 15
µg of Chromatin

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID

4087/4088

Swiss Prot

P84022/Q15796

Immunogen

This information is considered to be commercially sensitive.

Synonyms

CHTD8; JV18; JV18-1; LDS6; MADH2; MADR2; hMAD-2; hSMAD2; HSPC193; HsT17436; JV15-2; LDS1C; LDS3; MADH3; hMAD-3; hSMAD3; mad3

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immunoprecipitation of SMAD2/SMAD3 from 300 µg extracts of HeLa cells was performed using 0.7 µg of [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) at a dilution of 1:10000.




Western blot analysis of lysates from C2C12 cells using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) at 1:7000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10 s.




Western blot analysis of lysates from wild type (WT) and SMAD2 knockout (KO) HeLa cells using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) at 1:7000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10 s.


Validation Data



Western blot analysis of lysates from wild type (WT) and SMAD3 knockout (KO) 293T cells using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) at 1:7000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10 s.




Western blot analysis of lysates from wild type (WT) and SMAD2 knockout (KO) NIH/3T3 cells using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45 s.




Western blot analysis of various lysates using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20 s.


Validation Data




Chromatin immunoprecipitation was performed with 15 µg of cross-linked chromatin from HaCaT cells were treated with TGF-β (50 ng/mL,30 minutes), using 1 µg of [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.



Confocal imaging of HeLa cells (treated with hTGF-β3) and HeLa cells (untreated) using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434, dilution 1:1400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of NIH/3T3 cells using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434, dilution 1:1400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of PC-12 cells using [KO Validated] SMAD2/SMAD3 Rabbit mAb (A29434, dilution 1:1400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.