Leader in Biomolecular Solutions for Life Science

www.abclonal.com

MEK1/2 Rabbit mAb

Catalog No.: A29491 Recombinant

Basic Information

Observed MW
45 kDa

Calculated MW
41 kDa/43 kDa/44 kDa

Category
Primary antibody

Applications
WB,IF/ICC,ELISA

Cross-Reactivity
Human, Mouse, Rat, Monkey

CloneNo number
ARC4150

Background

The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development.

Recommended Dilutions

WB	1:5000 - 1:30000
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000)，a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID
5604/5605

Swiss Prot
P36507/Q02750

Immunogen
This information is considered to be commercially sensitive.

Synonyms

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃


Contact

400-999-6126


cn.market@abclonal.com.cn

www.abclonal.com.cn


Validation Data



Western blot analysis of various lysates using MEK1/2 Rabbit mAb (A29491) at 1:9000 dilution incubated overnight at 4℃.|Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.|Lysates/proteins: 25 µg per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit (RM00020).|Exposure time: 1 s.




Western blot analysis of various lysates using MEK1/2 Rabbit mAb (A29491) at 1:9000 dilution incubated overnight at 4℃.|Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.|Lysates/proteins: 25 µg per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit (RM00020).|Exposure time: 1 s.




Western blot analysis of various lysates using MEK1/2 Rabbit mAb (A29491) at 1:10000 dilution incubated overnight at 4℃.|Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.|Lysates/proteins: 25 µg per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit (RM00020).|Exposure time: 5 s.

Validation Data



Confocal imaging of HeLa cells using MEK1/2 Rabbit mAb (A29491, dilution 1:50) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of NIH/3T3 cells using MEK1/2 Rabbit mAb (A29491, dilution 1:50) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.