

COX2/PTGS2 Rabbit mAb

Catalog No.: A3560 Recombinant 33 Publications

Basic Information

Observed MW

74kDa

Calculated MW

69kDa

Category

Primary antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Clone/No. number

ARC0800

Background

Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis.

Recommended Dilutions

WB 1:1000 - 1:6000

IHC-P 1:200 - 1:2000

ELISA Recommended starting concentration is 1 μ g/mL.
 Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

5743

Swiss Prot

P35354

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

COX2; COX-2; PHS-2; PGG/HS; PGHS-2; hCox-2; GRIPGHS; COX2/PTGS2

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data

