# ATG16L1 Rabbit mAb

Catalog No.: A3637 Recombinant 1 Publications



### **Basic Information**

#### **Observed MW**

68kDa/72kDa

#### **Calculated MW**

68kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

#### CloneNo number

ARC0812

## **Background**

The protein encoded by this gene is part of a large protein complex that is necessary for autophagy, the major process by which intracellular components are targeted to lysosomes for degradation. Defects in this gene are a cause of susceptibility to inflammatory bowel disease type 10 (IBD10). Several transcript variants encoding different isoforms have been found for this gene.

### **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID

Swiss Prot

55054

Q676U5

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 6-105 of human ATG16L1 (Q676U5).

### **Synonyms**

IBD10; WDR30; APG16L; ATG16A; ATG16L; ATG16L1

### **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using ATG16L1 Rabbit mAb (A3637) at 1:1000 dilution

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.