

# iNOS Rabbit mAb

Catalog No.: A3774    Recombinant    51 Publications

## Basic Information

### Observed MW

131 kDa

### Calculated MW

131 kDa

### Category

Primary antibody

### Applications

WB, IF/ICC, ELISA

### Cross-Reactivity

Mouse, Rat

### Clone/No. number

ARC0832

## Background

Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission and antimicrobial and antitumoral activities. This gene encodes a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. Three related pseudogenes are located within the Smith-Magenis syndrome region on chromosome 17.

## Recommended Dilutions

**WB**                    1:1000 - 1:4000

**IF/ICC**                1:50 - 1:200

**ELISA**                Recommended starting concentration is 1  $\mu$ g/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

4843

### Swiss Prot

P35228

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

NOS; INOS; NOS2A; HEP-NOS; iNOS

## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data



Western blot analysis of lysates from RAW 264.7 cells using iNOS Rabbit mAb (A3774) at 1:1000 dilution incubated overnight at 4°C. Raw264.7 cells were treated with LPS (1 µg/ml) at 37°C for 8 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30 s.



Western blot analysis of lysates from NR8383 cells using iNOS Rabbit mAb (A3774) at 1:1000 dilution incubated at room temperature for 1.5 hours. NR8383 cells were treated with LPS (1 µg/ml) for 8 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5 s.



Confocal imaging of RAW 264.7 cells (treated with LPS) and RAW 264.7 cells(untreated) using iNOS Rabbit mAb (A3774, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.