# **NQO2** Rabbit mAb

Catalog No.: A3846 Recombinant



# **Basic Information**

#### **Observed MW**

26 kDa

### **Calculated MW**

26 kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Mouse, Rat

#### CloneNo number

ARC0846

# **Background**

This gene encodes a member of the thioredoxin family of enzymes. It is a cytosolic and ubiquitously expressed flavoprotein that catalyzes the two-electron reduction of quinone substrates and uses dihydronicotinamide riboside as a reducing coenzyme. Mutations in this gene have been associated with neurodegenerative diseases and several cancers. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 1:1000 - 1:4000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Swiss Prot Gene ID** 4835 P16083

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

QR2; DHQV; DIA6; NMOR2; NQO2

# **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn


# **Product Information**

**Purification Source** Isotype Rabbit IgG Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using NQO2 Rabbit mAb (A3846) at 1:1000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.