

# JAB1/CSN5/COPS5 Rabbit mAb

Catalog No.: A4087

Recombinant

2 Publications

## Basic Information

### Observed MW

35kDa

### Calculated MW

38kDa

### Category

Primary antibody

### Applications

WB, IHC-P, IF/ICC, ELISA, ChIP

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

ARC0895

## Background

The protein encoded by this gene is one of the eight subunits of COP9 signalosome, a highly conserved protein complex that functions as an important regulator in multiple signaling pathways. The structure and function of COP9 signalosome is similar to that of the 19S regulatory particle of 26S proteasome. COP9 signalosome has been shown to interact with SCF-type E3 ubiquitin ligases and act as a positive regulator of E3 ubiquitin ligases. This protein is reported to be involved in the degradation of cyclin-dependent kinase inhibitor CDKN1B/p27Kip1. It is also known to be an coactivator that increases the specificity of JUN/AP1 transcription factors.

## Recommended Dilutions

<b>WB</b>	1:500 - 1:2000
-----------	----------------

<b>IHC-P</b>	1:50 - 1:200
--------------	--------------

<b>IF/ICC</b>	1:50 - 1:200
---------------	--------------

<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
--------------	---

<b>ChIP</b>	5µg antibody for 10µg-15µg of Chromatin
-------------	---

## Immunogen Information

### Gene ID

10987

### Swiss Prot

Q92905

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

CSN5; JAB1; SGN5; MOV-34; JAB1/CSN5/COPS5

## Contact

		400-999-6126
--	--	--------------

		<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
--	--	--

		<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>
--	--	--

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data




Western blot analysis of various lysates using JAB1/CSN5/COP55 Rabbit mAb (A4087) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using JAB1/CSN5/COP55 antibody (A4087) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Immunofluorescence analysis of NIH-3T3 cells using JAB1/CSN5/COP55 Rabbit mAb (A4087) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of U-2 OS cells using JAB1/CSN5/COP55 Rabbit mAb (A4087) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using JAB1/CSN5/COP55 Rabbit mAb (A4087) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


## Validation Data




Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using JAB1/CSN5/COPS5 Rabbit mAb (A4087) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using JAB1/CSN5/COPS5 Rabbit mAb (A4087) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using JAB1/CSN5/COPS5 Rabbit mAb (A4087) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using JAB1/CSN5/COPS5 Rabbit mAb (A4087) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis tissue using JAB1/CSN5/COPS5 Rabbit mAb (A4087) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.