Leader in Biomolecular Solutions for Life Science



Catalog No.: A4151 5 Publications



Basic Information

Observed MW 48kDa

Calculated MW 48kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat

Background

This gene product is a member of the transcriptional enhancer factor (TEF) family of transcription factors, which contain the TEA/ATTS DNA-binding domain. It is preferentially expressed in the skeletal muscle, and binds to the M-CAT regulatory element found in promoters of muscle-specific genes to direct their gene expression. Alternatively spliced transcripts encoding distinct isoforms, some of which are translated through the use of a non-AUG (UUG) initiation codon, have been described for this gene.

Recommended Dilutions

WB	1:200 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID 7004

Swiss Prot Q15561

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TEF3; RTEF1; TEF-3; EFTR-2; TEFR-1; TCF13L1; hRTEF-1B; TEAD4

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using TEAD4 Rabbit pAb (A4151) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.