

[KO Validated] SMARCC1/BAF155 Rabbit mAb

Catalog No.: A4275 KO Validated Recombinant

Basic Information

Observed MW

155kDa/

Calculated MW

123kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA,ChIP

Cross-Reactivity

Human, Mouse

CloneNo number

ARC0948

Background

The protein encoded by this gene is a member of the SWI/SNF family of proteins, whose members display helicase and ATPase activities and which are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI and contains a predicted leucine zipper motif typical of many transcription factors.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

ELISA Recommended starting

concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay

requirements.

ChIP 5μg antibody for

10μg-15μg of Chromatin

Immunogen Information

Gene ID Swiss Prot 6599 Q92922

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

HYC5; Rsc8; SRG3; SWI3; BAF155; CRACC1; 55

Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
<u> </u>	I	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using SMARCC1/BAF155 Rabbit mAb (A4275) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3min.




Western blot analysis of lysates from wild type (WT) and SMARCC1/BAF155 knockout (KO) 293T cells using [KO Validated] SMARCC1/BAF155 Rabbit mAb (A4275) at 1:1000 dilution incubated overnight at 4° C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 30s.



Chromatin immunoprecipitation analysis of extracts of MCF7 beta-estradiol cells, using SMARCC1/BAF155 antibody (A4275) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Immunofluorescence analysis of NIH-3T3 cells using SMARCC1/BAF155 Rabbit mAb (A4275) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.