[KO Validated] MTA2 Rabbit mAb

www.abclonal.com

ABclonal

Catalog No.: A4624 KO Validated Recombinant

Basic Information

Observed MW

75kDa

Calculated MW

75kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,ELISA,ChIP

Cross-Reactivity

Human

CloneNo number

ARC1056

Background

This gene encodes a protein that has been identified as a component of NuRD, a nucleosome remodeling deacetylase complex identified in the nucleus of human cells. It shows a very broad expression pattern and is strongly expressed in many tissues. It may represent one member of a small gene family that encode different but related proteins involved either directly or indirectly in transcriptional regulation. Their indirect effects on transcriptional regulation may include chromatin remodeling. It is closely related to another member of this family, a protein that has been correlated with the metastatic potential of certain carcinomas. These two proteins are so closely related that they share the same types of domains. These domains include two DNA binding domains, a dimerization domain, and a domain commonly found in proteins that methylate DNA. One of the proteins known to be a target protein for this gene product is p53. Deacetylation of p53 is correlated with a loss of growth inhibition in transformed cells supporting a connection between these gene family members and metastasis.

Recommended Dilutions

WB 1:500 - 1:1000

1:50 - 1:200 **IHC-P**

1:50 - 1:200 IF/ICC

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on vour specific assav

requirements.

ChIP 5µg antibody for

10μg-15μg of Chromatin

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Immunogen Information

Gene ID Swiss Prot 9219 094776

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PID; MTA1L1; A2


Product Information

Source Isotype **Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from wild type (WT) and MTA2 knockout (KO) 293T cells,using [KO Validated] MTA2 Rabbit mAb (A4624) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Western blot analysis of various lysates using [KO Validated] MTA2 Rabbit mAb (A4624) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 3min.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using [KO Validated] MTA2 Rabbit mAb (A4624) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Immunofluorescence analysis of U-2 OS cells using [KO Validated] MTA2 Rabbit mAb (A4624) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffinembedded Human colon using [KO Validated] MTA2 Rabbit mAb (A4624) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.