# PER2 Rabbit mAb

Catalog No.: A5107 Recombinant 4 Publications



## **Basic Information**

### **Observed MW**

140kDa

#### **Calculated MW**

137kDa

## Category

Primary antibody

### **Applications**

ELISA, WB, ChIP

#### **Cross-Reactivity**

Human

### CloneNo number

ARC1134

## **Background**

This gene is a member of the Period family of genes and is expressed in a circadian pattern in the suprachiasmatic nucleus, the primary circadian pacemaker in the mammalian brain. Genes in this family encode components of the circadian rhythms of locomotor activity, metabolism, and behavior. This gene is upregulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback loop using PER/CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene may increase the risk of getting certain cancers and have been linked to sleep disorders.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ChiP** 5μg antibody for 10μg-15μg of Chromatin

## **Immunogen Information**

**Gene ID**8864
Swiss Prot
015055

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human PER2 (O15055).

## **Synonyms**

FASPS; FASPS1; PER2

## **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.




Western blot analysis of various lysates using PER2 Rabbit mAb (A5107) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 3min.



Chromatin immunoprecipitation analysis of extracts of MCF7 cells, using PER2 antibody (A5107) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.