# PIAS1 Rabbit pAb

Catalog No.: A5729



### **Basic Information**

#### **Observed MW**

Refer to figures

### **Calculated MW**

72kDa

### Category

Primary antibody

### **Applications**

ELISA,IF/ICC

### **Cross-Reactivity**

Human

## **Background**

This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins function as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. This protein plays a central role as a transcriptional coregulator of numerous cellular pathways including the STAT1 and nuclear factor kappaB pathways. Alternate splicing results in multiple transcript variants.

## **Recommended Dilutions**

IF/ICC

1:50 - 1:200

## **Immunogen Information**

Gene ID 8554 **Swiss Prot** 

075925

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 422-651 of human PIAS1 (NP $_0$ 57250.1).

### **Synonyms**

GBP; ZMIZ3; DDXBP1; GU/RH-II; PIAS1

### **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification


### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Immunofluorescence analysis of GFP-RNF168 transgenic U2OS cells using PIAS1 antibody (A5729). Green: GFP-RNF168 fusion protein expression for DNA damage marker. Blue: DAPI for nuclear staining. RNF168 (GFP) can be used to mark cells damaged by UV-A laser for they always gather around DNA damage region.



Immunofluorescence analysis of U2OS cells using PIAS1 antibody (A5729). Blue: DAPI for nuclear staining.