FADD Rabbit pAb

Catalog No.: A5819 10 Publications



Basic Information

Observed MW

26kDa

Calculated MW

23kDa

Category

Primary antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.

Recommended Dilutions

WB 1:500 - 1:2000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot8772
Q13158

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GIG3; IMD90; MORT1; FADD

Contact

| a | 400-999-6126 |
|-------------------|---------------------------|
| × | cn.market@abclonal.com.cn |
| $\overline{\Box}$ | www.ahclonal.com.cn |


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using FADD Rabbit pAb (A5819) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).



Immunoprecipitation analysis of 200 μ g extracts of HeLa cells using 3 μ g FADD antibody (A5819). Western blot was performed from the immunoprecipitate using FADD antibody (A5819) at a dilution of 1:1000.