

STRAP Rabbit pAb

Catalog No.: A5964

1 Publications

Basic Information

Observed MW

39kDa

Calculated MW

38kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables RNA binding activity. Involved in maintenance of gastrointestinal epithelium; negative regulation of transforming growth factor beta receptor signaling pathway; and spliceosomal snRNP assembly. Located in cytosol. Part of SMN complex. Implicated in adenocarcinoma; colorectal carcinoma; large cell carcinoma; lung carcinoma; and squamous cell neoplasm.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

11171

Swiss Prot

Q9Y3F4

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

MAWD; PT-WD; UNRIP; STRAP

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of various lysates using STRAP Rabbit pAb (A5964) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates / proteins: 25 µg per lane.
Blocking buffer: 3 % nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5s.