# P2RX4 Rabbit pAb

Catalog No.: A6682 2 Publications



# **Basic Information**

#### **Observed MW**

62kDa

### **Calculated MW**

43kDa

# Category

Primary antibody

### **Applications**

ELISA,WB,IF/ICC

#### **Cross-Reactivity**

Human, Mouse

# **Background**

The product of this gene belongs to the family of purinoceptors for ATP. This receptor functions as a ligand-gated ion channel with high calcium permeability. The main pharmacological distinction between the members of the purinoceptor family is the relative sensitivity to the antagonists suramin and PPADS. The product of this gene has the lowest sensitivity for these antagonists. Multiple alternatively spliced transcript variants, some protein-coding and some not protein-coding, have been found for this gene.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:10 - 1:100

# **Immunogen Information**

**Gene ID Swiss Prot** 5025 Q99571

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 55-338 of human P2RX4 (NP\_002551.2).

### **Synonyms**

P2X4; P2X4R; P2RX4

# **Contact**

	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using P2RX4 antibody (A6682) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunofluorescence analysis of MCF7 cells using P2RX4 antibody (A6682). Blue: DAPI for nuclear staining.