

# Acetyl-Histone H3-K18 Rabbit pAb

Catalog No.: A7257 **16 Publications**

## Basic Information

### Observed MW

17kDa

### Calculated MW

15kDa

### Category

Primary antibody

### Applications

WB,DB,IHC-P,IF/ICC,ELISA,CHIP

### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**DB** 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

**CHIP** 5µg antibody for 5µg-10µg of Chromatin

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J; Acetyl-Histone H3-K18

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




| [www.abclonal.com.cn](http://www.abclonal.com.cn)

---


## Validation Data






Western blot analysis of lysates from HeLa cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at 1:1000 dilution. HeLa cells were treated with TSA (1  $\mu$ M) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 0.5s.



Chromatin immunoprecipitation analysis of extracts of HCT116 cells, using Acetyl-Histone H3-K18 Rabbit pAb antibody (A7257) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Chromatin immunoprecipitation analysis of extracts of HCT116 cells, using Acetyl-Histone H3-K18 Rabbit pAb antibody (A7257) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



## Validation Data


---

Dot-blot analysis of all sorts of methylation peptides using Acetyl-Histone H3-K18 antibody (A7257) at 1:1000 dilution.




Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100. NIH/3T3 cells were treated with TSA (1  $\mu$ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of C6 cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100. C6 cells were treated with TSA (1  $\mu$ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded Rat brain using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunofluorescence analysis of HeLa cells using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100. HeLa cells were treated with TSA (1  $\mu$ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Human gastric cancer using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Acetyl-Histone H3-K18 Rabbit pAb (A7257) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.