NOX1 Rabbit pAb

Catalog No.: A8527 9 Publications



Basic Information

Observed MW

65kDa

Calculated MW

65kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the NADPH oxidase family of enzymes responsible for the catalytic one-electron transfer of oxygen to generate superoxide or hydrogen peroxide. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Swiss Prot Gene ID 27035 Q9Y5S8

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

MOX1; NOH1; NOH-1; GP91-2; NOH-1L; NOX1

Contact

| a | 400-999-6126 |
|----------|---------------------------|
| \sim | cn.market@abclonal.com.cn |
| ⊙ | www.abclonal.com.cn |


Product Information

Purification Source Isotype Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using NOX1 Rabbit pAb (A8527) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (ASO14) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.