

PCNA Mouse mAb

Catalog No.: AC058

Basic Information

Observed MW

36kDa

Calculated MW

29kDa

Category

Loading control antibody

Applications

WB,IF/ICC,IF-P,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

AMC0704

Background

Enables estrogen receptor binding activity. Involved in several processes, including estrous cycle; liver regeneration; and response to dexamethasone. Located in nucleus. Biomarker of several diseases, including brain ischemia (multiple); mucositis; obstructive jaundice; osteoporosis; and sciatic neuropathy. Human ortholog(s) of this gene implicated in stomach cancer. Orthologous to human PCNA (proliferating cell nuclear antigen).

Recommended Dilutions

WB	1:1000 - 1:10000
IF/ICC	1:200 - 1:800
IF-P	1:200 - 1:800
IHC-P	1:2000-1:8000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Immunogen Information

Gene ID

25737

Swiss Prot

P04961

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

PCNAR; PcnA/cyclin

Product Information

Source

Mouse

Isotype

IgG2a, κ


Purification

Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.02% sodium azide, pH7.2.

Validation Data



Western blot analysis of various lysates using PCNA Mouse mAb (AC058) at 1:5000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Confocal imaging of HeLa cells using PCNA Mouse mAb (AC058, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of NIH/3T3 cells using PCNA Mouse mAb (AC058, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of paraffin-embedded Rat small intestine tissue using PCNA Mouse mAb (AC058, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffin-embedded Mouse skin tissue using PCNA Mouse mAb (AC058) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using PCNA Mouse mAb (AC058) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon tissue using PCNA Mouse mAb (AC058) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using PCNA Mouse mAb (AC058) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.