

HRP-conjugated Mouse anti HA-Tag mAb

Catalog No.: AE025 9 Publications

Basic Information

Observed MW

53kDa(CD95)/65kDa(CD2)/53kDa(SCRN1)
)/26kDa(HA(N
))/42kDa(WDR77)/45kDa(NR3C2)

Calculated MW

Category

Tag antibody

Applications

WB,ELISA

Cross-Reactivity

Species independent

Clone/No. number

AMC0557

Conjugate

HRP

Recommended Dilutions

WB 1:10000 - 1:20000

ELISA Recommended starting concentration is 1 μ g/mL.
 Please optimize the concentration based on your specific assay requirements.

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

HA;HA tag;HA-tag

Product Information

Source

Mouse

Isotype

IgG1,Kappa

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Validation Data



Western blot analysis of extracts of normal 293T cells, 293T transfected with SCRN1 Protein , using HRP-conjugated Mouse anti HA-Tag mAb (AE025) at 1:5000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Western blot analysis of extracts of normal 293T and 293F cells, 293T transfected with CD2 Protein and HA-N Protein ,293F transfected with CD95-N Protein using HRP-conjugated Mouse anti HA-Tag mAb (AE025) at 1:5000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3s.



Western blot analysis of extracts of normal 293T and 293F cells, 293T transfected with NR3C2-C Protein,293F transfected with WDR77-C Protein using HRP-conjugated Mouse anti HA-Tag mAb (AE025) at 1:5000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.