

Myc-Tag Rabbit mAb (C-terminal)

Catalog No.: AE070 **Recombinant** **75 Publications**

Basic Information

Observed MW

58 kDa (NLK transfected)

Calculated MW

Category

Tag antibody

Applications

WB, IF/ICC, IP, ELISA

Cross-Reactivity

Species independent

CloneNo number

ARC5004-12

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB 1:10000 - 1:120000**IF/ICC** 1:200 - 1:1000**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Myc; Myc tag; Myc-tag; Myc-Tag

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Immunoprecipitation analysis of 200ug extracts of 293T cells transfected with NLK expression vector containing a Myc-Tag Rabbit mAb (C-terminal) (AE070 1:100). Western blot was performed from the immunoprecipitate using Myc-Tag Rabbit mAb (C-terminal) at 1:1000 dilution.



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with using NLK-Myc-Tag Rabbit mAb (C-terminal) (AE070) at 1:120000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10 s.



Confocal imaging of HeLa cells transfected with H2B-EGFP-Myc(C) using Myc-Tag Rabbit mAb (C-terminal) (AE070, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.