

Rabbit anti GST-Tag mAb

Catalog No.: AE077 8 Publications

Basic Information

Observed MW

35-50 kDa

Calculated MW

26kDa

Category

Tag antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Species independent

Clone/No number

ARC50738

Background

Glutathione S-transferases (GSTs), previously known as ligandins, comprise a family of eukaryotic and prokaryotic phase II metabolic isozymes best known for their ability to catalyze the conjugation of the reduced form of glutathione (GSH) to xenobiotic substrates for the purpose of detoxification. The GST family consists of three superfamilies: the cytosolic, mitochondrial, and microsomal—also known as MAPEG—proteins. Members of the GST superfamily are extremely diverse in amino acid sequence, and a large fraction of the sequences deposited in public databases are of unknown function. The Enzyme Function Initiative (EFI) is using GSTs as a model superfamily to identify new GST functions. A GST-tag is often used to separate and purify proteins that contain the GST-fusion protein. The tag is 220 amino acids (roughly 26 kDa) in size, which, compared to tags such as the Myc-tag or the FLAG-tag, is quite large. It can be fused to either the N-terminus or C-terminus of a protein. However, many commercially available sources of GST-tagged plasmids include a thrombin domain for cleavage of the GST tag during protein purification.

Recommended Dilutions

| | |
|-------|---|
| WB | 1:1000 - 1:6000 |
| IP | 3μg antibody for 1μg extracts of recombinant protein |
| ELISA | Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements. |

Immunogen Information

Gene ID

Swiss Prot
P08515

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GST;GST tag;GST-tag

Contact

| | |
|--|--|
|  | 400-999-6126 |
|  | cn.market@abclonal.com.cn |
|  | www.abclonal.com.cn |

Product Information


| Source | Isotype | Purification |
|--------|---------|-----------------------|
| Rabbit | IgG | Affinity purification |

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Immunoprecipitation of GST-Tag from 300 μ g extracts of 293F cells transfected with GST-Tag was performed using 2 μ g of Rabbit anti GST-Tag mAb (AE077). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10 % of the total input. Western blot analysis of immunoprecipitates was conducted using Rabbit anti GST-Tag mAb (AE077) at a dilution of 1:2000.

