

# Rabbit anti V5-Tag mAb

Catalog No.: AE089

3 Publications

## Basic Information

**Observed MW**

40kDa

**Calculated MW****Category**

Tag antibody

**Applications**

WB, IF/ICC, ELISA, ChIP

**Cross-Reactivity**

Species independent

**CloneNo number**

ARC50445

## Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

## Recommended Dilutions

**WB** 1:2000 - 1:6000**IF/ICC** 1:50 - 1:200**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.**ChIP** 5µg antibody for 10µg-15µg of Chromatin

## Immunogen Information


**Gene ID****Swiss Prot****Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

V5;V5 tag;V5-tag

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**




Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Immunofluorescence analysis of 293T cells transfected with V5(mCherry) protein and untreated 293T cells using Rabbit anti V5-Tag mAb (AE089) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.