

# PE Mouse anti DDDDK-Tag mAb (C-terminal)

Catalog No.: AE116

2 Publications

## Basic Information

### Observed MW

### Calculated MW

### Category

Tag antibody

### Applications

FC (intra)

### Cross-Reactivity

Species independent

### CloneNo number

AMC0382

### Conjugate

PE. Ex:565nm. Em:574nm.

## Background

FLAG-tag, or FLAG octapeptide, or FLAG epitope, is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. It has been used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits, because its mild purification procedure tends not to disrupt such complexes. It has been used to obtain proteins of sufficient purity and quality to carry out 3D structure determination by x-ray crystallography. A FLAG-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against a given protein, adding a FLAG-tag to a protein allows the protein to be studied with an antibody against the FLAG sequence. Examples are cellular localization studies by immunofluorescence or detection by SDS PAGE protein electrophoresis and Western blotting.

## Recommended Dilutions

**FC (intra)** 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Immunogen Information

### Gene ID

### Swiss Prot

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

DDDDK;DDDDK tag;DDDDK-tag

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Mouse

### Isotype

IgG1, Kappa

### Purification


Affinity purification

### Storage


Store at 2-8°C. Avoid freeze / thaw cycles.

Buffer: PBS, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

# Validation Data



Flow cytometry: 1X10<sup>6</sup> CHO cells (negative control) and CHO(Transfection, right) cells were intracellularly-stained with PE Mouse anti DDDDK-Tag mAb(AE116, 5  $\mu$ l/Test, orange line) or PE Mouse IgG isotype control (5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10<sup>6</sup> CHO(Transfection) cells were intracellularly-stained with PE Mouse IgG isotype control (5  $\mu$ l/Test, left) or PE Mouse anti DDDDK-Tag mAb(AE116, 5  $\mu$ l/Test, right).