

Rabbit anti mCherry-Tag mAb

Catalog No.: AE127

1 Publications

Basic Information

Observed MW

33kDa

Calculated MW**Category**

Tag antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Species independent

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB 1:10000 - 1:40000**IP** 0.5µg-4µg antibody for
200µg-600µg extracts of
whole cells**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID**Swiss Prot****Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

mCherry; mCherry tag; mCherry-tag

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Immunoprecipitation of CT55-mCherry from 300 µg extracts of 293F cells transfected with a CT55 expression vector containing a single C-terminal mCherry-Tag was performed using 3 µg of mCherry-Tag Rabbit mAb (AE127). Rabbit IgG isotype control (AC042) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10 % of the total input. Western blot analysis of immunoprecipitates was conducted using mCherry-Tag Rabbit mAb (AE127) at a dilution of 1:6000.



Western blot analysis of lysates from wild type (WT) and 293F cells transfected with mCherry using Rabbit anti mCherry-Tag mAb (AE127) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 20 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45s.