

# Phospho-CDK1-Y15 Rabbit pAb

Catalog No.: AP0016 **21 Publications**

## Basic Information

### Observed MW

34kDa

### Calculated MW

34kDa

### Category

Primary antibody

### Applications

WB,IP,IF-P,IHC-P,ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

<b>WB</b>	1:500 - 1:1000
<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
<b>IF-P</b>	1:50 - 1:200
<b>IHC-P</b>	1:50 - 1:200
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Immunogen Information

### Gene ID

983

### Swiss Prot

P06493

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

CDC2; CDC28A; P34CDC2; Phospho-CDK1-Y15

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Immunoprecipitation analysis of 200 µg extracts of HT-29 cells, using 3 µg Phospho-CDK1-Y15 pAb (AP0016). Western blot was performed from the immunoprecipitate using Phospho-CDK1-Y15 pAb (AP0016) at a dilution of 1:1000. HT-29 cells were treated with Serum-starvation overnight at 37°C.



Western blot analysis of lysates from HeLa cells, using Phospho-CDK1-Y15 Rabbit pAb (A0220). HeLa cells were treated with nocodazole (50 ng/mL) at 37°C for 20 hours or Hydroxyurea (4 mM) at 37°C for 20 hours.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of various lysates using Phospho-CDK1-Y15 Rabbit pAb (AP0016) at 1:1000 dilution.

HeLa and NIH/3T3 cells were treated with nocodazole (50 ng/ml) at 37°C for 20 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.


## Validation Data




Western blot analysis of lysates from C6 cells, using Phospho-CDK1-Y15 Rabbit pAb (AP0016) at 1:1000 dilution. C6 cells were treated with Nocodazole (50 ng/ml) at 37°C for 20 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.




Immunofluorescence analysis of paraffin-embedded human breast cancer using Phospho-CDK1-Y15 Rabbit pAb (AP0016) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Human colon using Phospho-CDK1-Y15 Rabbit pAb (AP0016) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver using Phospho-CDK1-Y15 Rabbit pAb (AP0016) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Phospho-CDK1-Y15 Rabbit pAb (AP0016) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.