

# Phospho-ERK1-Y204 Rabbit pAb

Catalog No.: AP0235

12 Publications

## Basic Information

### Observed MW

42kDa/44kDa

### Calculated MW

43kDa

### Category

Primary antibody

### Applications

WB, IHC-P, IF/ICC, IP, ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described.

## Recommended Dilutions

**WB** 1:500 - 1:1000**IHC-P** 1:50 - 1:100**IF/ICC** 1:100 - 1:200**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Immunogen Information

### Gene ID

5595

### Swiss Prot

P27361

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; Phospho-ERK1-Y204

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)


## Validation Data




Immunoprecipitation analysis of 200 µg extracts of Jurkat cells, using 3 µg Phospho-ERK1-Y204 Rabbit pAb (AP0235). Western blot was performed from the immunoprecipitate using Phospho-ERK1-Y204 Rabbit pAb (AP0235) at a dilution of 1:1000. Jurkat cells were treated with PMA/TPA (200 nM) at 37°C for 10 minutes.




Western blot analysis of lysates from NIH/3T3 cells using Phospho-ERK1-Y204 Rabbit pAb (AP0235) at 1:1000 dilution incubated overnight at 4°C. NIH/3T3 cells were treated with PMA/TPA (200 nM) at 37°C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Immunofluorescence analysis of methanol-fixed HeLa cells showing centrosome and nuclear staining using Phospho-ERK1-Y204 Rabbit pAb (AP0235).




Immunohistochemistry analysis of paraffin-embedded Rat liver using Phospho-ERK1-Y204 Rabbit pAb (AP0235) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human uterus using Phospho-ERK1-Y204 Rabbit pAb (AP0235) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.

## Validation Data

---



Immunohistochemistry analysis of paraffin-embedded Mouse heart using Phospho-ERK1-Y204 Rabbit pAb (AP0235) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.