

# Phospho-ZAP70-Y319 Rabbit pAb

**Catalog No.: AP0467**

## Basic Information

**Observed MW**

70kDa

**Calculated MW**

70kDa

**Category**

Primary antibody

**Applications**

WB,IP,ELISA

**Cross-Reactivity**

Human

## Background

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.

## Recommended Dilutions

**WB** 1:500 - 1:2000**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Immunogen Information

**Gene ID**

7535

**Swiss Prot**

P43403

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

SRK; STD; TZK; STCD; IMD48; ADMIO2; ZAP-70; Phospho-ZAP70-Y319

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# Validation Data



Immunoprecipitation analysis of 200 µg extracts of Jurkat cells, using 3 µg Phospho-ZAP70-Y319 pAb (AP0467). Western blot was performed from the immunoprecipitate using Phospho-ZAP70-Y319 pAb (AP0467) at a dilution of 1:1000. Jurkat cells were treated with Hydrogen Peroxide (2 mM) at 37°C for 2 minutes after serum-starvation overnight.