

Phospho-ERK1-T202/Y204 + ERK2-T185/Y187 Rabbit pAb

Catalog No.: AP0472 138 Publications

Basic Information

Observed MW

44kDa/ 42kDa/

Calculated MW

36kDa/41kDa/38kDa/40kDa/43kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. One study also suggests that this protein acts as a transcriptional repressor independent of its kinase activity. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Two alternatively spliced transcript variants encoding the same protein, but differing in the UTRs, have been reported for this gene. [provided by RefSeq, Jan 2014]

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 5594/5595 P28482/P27361

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

MAPK1/MAPK3; Phospho-ERK1-T202/Y204 + ERK2-T185/Y187

Contact

a	400-999-6126
\sim	cn.market@abclonal.com.cn
⊙	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from C6 cells using Phospho-ERK1-T202/Y204 + ERK2-T185/Y187 Rabbit pAb (AP0472) at 1:1000 dilution incubated overnight at 4° C. C6 cells were treated with PMA(200 nM) at 37° C for 10 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.



Western blot analysis of lysates from NIH/3T3 cells using Phospho-ERK1-T202/Y204 + ERK2-T185/Y187 Rabbit pAb (AP0472) at 1:1000 dilution. NIH/3T3 cells were treated with PMA/TPA (200 nM) at 37° C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.