

Phospho-PKA C-alpha (PRKACA)-S339 Rabbit pAb

Catalog No.: AP0558

Basic Information

Observed MW

42kDa

Calculated MW

41kDa

Category

Primary antibody

Applications

WB, IP, ELISA

Cross-Reactivity

Human

Background

This gene encodes one of the catalytic subunits of protein kinase A, which exists as a tetrameric holoenzyme with two regulatory subunits and two catalytic subunits, in its inactive form. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. cAMP-dependent phosphorylation of proteins by protein kinase A is important to many cellular processes, including differentiation, proliferation, and apoptosis. Constitutive activation of this gene caused either by somatic mutations, or genomic duplications of regions that include this gene, have been associated with hyperplasias and adenomas of the adrenal cortex and are linked to corticotropin-independent Cushing's syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. Tissue-specific isoforms that differ at the N-terminus have been described, and these isoforms may differ in the post-translational modifications that occur at the N-terminus of some isoforms.

Recommended Dilutions

WB 1:500 - 1:2000**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

5566

Swiss Prot

P17612

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

CAFD1; PKACA; PPNAD4; Phospho-PKA C-alpha (PRKACA)-S339

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Immunoprecipitation analysis of 200 µg extracts of HeLa cells, using 3 µg Phospho-PKA C-alpha (PRKACA)-S339 pAb (AP0558). Western blot was performed from the immunoprecipitate using Phospho-PKA C-alpha (PRKACA)-S339 pAb (AP0558) at a dilution of 1:1000. HeLa cells were treated with EGF (100 ng/mL) at 37°C for 30 minutes after serum-starvation overnight.



Western blot analysis of lysates from HeLa and 293 cells, using Phospho-PKA C-alpha (PRKACA)-S339 Rabbit pAb (AP0558) at 1:1000 dilution. HeLa cells were treated with EGF (100ng/mL) for 30 minutes after serum-starvation overnight. 293T cells were treated with Insulin (100nM) for 10 minutes after serum-starvation overnight. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.