

Phospho-IRE1-S724 Rabbit pAb

Catalog No.: AP0878SP **28 Publications**

Basic Information

Observed MW

130 kDa

Calculated MW

110 kDa

Category

Primary antibody

Applications

WB,IF/ICC,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes the transmembrane protein kinase inositol-requiring enzyme 1. The encoded protein contains two functional catalytic domains, a serine/threonine-protein kinase domain and an endoribonuclease domain. This protein functions as a sensor of unfolded proteins in the endoplasmic reticulum (ER) and triggers an intracellular signaling pathway termed the unfolded protein response (UPR). The UPR is an ER stress response that is conserved from yeast to mammals and activates genes involved in degrading misfolded proteins, regulating protein synthesis and activating molecular chaperones. This protein specifically mediates the splicing and activation of the stress response transcription factor X-box binding protein 1.

Recommended Dilutions

WB 1:1000 - 1:2000

IF/ICC 1:100 - 1:200

IHC-P 1:4000 - 1:16000

ELISA

Recommended starting concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ($\geq 1:10000$) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID

2081

Swiss Prot

O75460

Immunogen

This information is considered to be commercially sensitive.

Synonyms

IRE1; IRE1P; IRE1a; hIRE1p; Phospho-IRE1-S724

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide.

May contain 0.05% BSA as specified on the Certificate of Analysis.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Validation Data




Western blot analysis of various lysates using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at 1:1000 dilution incubated overnight at 4°C. HeLa cells treated with CA (100 nM) at 37°C for 30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20 s.




Western blot analysis of various lysates using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at 1:1000 dilution incubated overnight at 4°C. C6 cells treated with CA (50 nM) at 37°C for 30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20 s.



Immunofluorescence analysis of HeLa cells (treated with CA) and HeLa cells (untreated) using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells (treated with CA) and NIH/3T3 cells (untreated) using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of C6 cells (treated with CA) and C6 cells (untreated) using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue, untreated (left) and lambda phosphatase-treated (right), tissue using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon tissue, untreated (left) and lambda phosphatase-treated (right), tissue using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue, untreated (left) and lambda phosphatase-treated (right), tissue using Phospho-IRE1-S724 Rabbit pAb (AP0878SP) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.