Phospho-GSK3β-S9 Rabbit mAb

www.abclonal.com

ABclonal

Catalog No.: AP1088 Recombinant 19 Publications

Basic Information

Observed MW

47 kDa

Calculated MW

47 kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human

CloneNo number

ARC0069

Background

The protein encoded by this gene is a serine-threonine kinase belonging to the glycogen synthase kinase subfamily. It is a negative regulator of glucose homeostasis and is involved in energy metabolism, inflammation, ER-stress, mitochondrial dysfunction, and apoptotic pathways. Defects in this gene have been associated with Parkinson disease and Alzheimer disease.

Recommended Dilutions

WB 1:1000 - 1:6000

1:50 - 1:200 **IHC-P**

Recommended starting **ELISA** concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 2932 P49841

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

GSK3B; gsk-3β; Phospho-GSK3β-S9

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
$\overline{\alpha}$	www.abclonal.com.cn


Product Information

Source Isotype **Purification** lgG Rabbit Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from HeLa cells using Phospho-GSK3 β -S9 Rabbit mAb (AP1088) at 1:3000 dilution incubated at room temperature for 1.5 hours. HeLa cells were treated with Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 20s.



Immunohistochemistry analysis of paraffinembedded Human colon carcinoma tissue using Phospho-GSK3 β -S9 Rabbit mAb (AP1088) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human thyroid cancer tissue using Phospho-GSK3 β -S9 Rabbit mAb (AP1088) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human colon tissue using Phospho-GSK3 β -S9 Rabbit mAb (AP1088) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.