# Phospho-STK11-S428 Rabbit pAb

Catalog No.: AP1108



### **Basic Information**

#### **Observed MW**

65kDa

#### **Calculated MW**

49kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a serine/threonine kinase that regulates cell polarity and energy metabolism and functions as a tumor suppressor. Mutations in this gene have been associated with the autosomal dominant Peutz-Jeghers syndrome, as well as with skin, pancreatic, and testicular cancers.

## **Recommended Dilutions**

WB

1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

Gene ID 6794 Swiss Prot Q15831

## Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

PJS; LKB1; hLKB1; Phospho-STK11-S428

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

### **Validation Data**



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-STK11-S428 Rabbit pAb (AP1108) at 1:1000 dilution or LKB1 antibody (A2122). NIH/3T3 cells were treated with PMA/TPA (200 nM) at  $37^{\circ}$ C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins:  $25\mu g$  per lane. Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.