

Phospho-RIPK1/RIP-S166 Rabbit pAb

Catalog No.: AP1230

5 Publications

Basic Information

Observed MW

75kDa

Calculated MW

76kDa

Category

Primary antibody

Applications

ELISA, WB

Cross-Reactivity

Human, Mouse

Background

This gene encodes a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases. The encoded protein plays a role in inflammation and cell death in response to tissue damage, pathogen recognition, and as part of developmental regulation. RIPK1/RIPK3 kinase-mediated necrosis is referred to as necroptosis. Genetic disruption of this gene in mice results in death shortly after birth.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

8737

Swiss Prot

Q13546

Immunogen

A synthetic phosphorylated peptide around S166 of human RIPK1/RIP (NP_003795.2).

Synonyms

RIP; RIP1; AIEFL; IMD57; RIP-1; Phospho-RIPK1/RIP-S166

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of HT-29, using Phospho-RIPK1/RIP-S166 Rabbit pAb (AP1230) at 1:800 dilution. HT-29 cells were treated with Z-VAD (50uM), human TNF-α (50ng/ml) and SM-164 (200nM) at 37°C for 5 hour.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.