

Phospho-MAP2K4-S257 Rabbit mAb

Catalog No.: AP1491 **Recombinant**

Basic Information

Observed MW

44kDa

Calculated MW

44kDa

Category

Primary antibody

Applications

WB, IHC-P, IF/ICC, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC66577

Background

This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:1000 - 1:5000**IHC-P** 1:50 - 1:200**IF/ICC** 1:50 - 1:200**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

6416

Swiss Prot

P45985

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

JNKK; MEK4; MKK4; SEK1; SKK1; JNKK1; SERK1; MAPKK4; PRKMK4; SAPKK1; SAPKK-1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn


Validation Data




Immunoprecipitation of Phospho-MAP2K4-S257 from 200 µg extracts of HeLa cells treated by UV (100 mJ/cm², 4 hours) was performed using 0.5 µg of Phospho-MAP2K4-S257 Rabbit mAb (AP1491). Rabbit IgG isotype control (AC042) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:1000.




Western blot analysis of lysates from C6 cells using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at 1:2600 dilution incubated overnight at 4°C. C6 cells were treated by UV (90mJ/cm²) at room temperature and recovered for 2 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.



Confocal imaging of HeLa cells (treated with UV) and HeLa cells (untreated) using Phospho-MAP2K4-S257 Rabbit mAb (AP1491, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue, Untreated (left) and lambda phosphatase-treated (right), using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue, Untreated (left) and lambda phosphatase-treated (right), using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Rat spleen tissue, Untreated (left) and lambda phosphatase-treated (right), using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Phospho-MAP2K4-S257 Rabbit mAb (AP1491) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.