

# HRP-conjugated Goat anti-Rat IgG (H+L)

Catalog No.: AS028 33 Publications

## Basic Information

### Observed MW

### Calculated MW

### Category

Secondary antibody

### Applications

WB,IHC-P

### Cross-Reactivity

Rat

### Conjugate

HRP

## Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

## Recommended Dilutions

WB 1:5000 - 1:10000

IHC-P 1:500 - 1:2000

## Immunogen Information

### Gene ID

### Swiss Prot

### Immunogen

This information is considered to be commercially sensitive.

### Synonyms

## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>

## Product Information

### Source

Goat

### Isotype

Horseradish peroxidase conjugated IgG


### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.  
Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from Rat IgG using HRP Goat Anti-Rat IgG (H+L) (AS028) at 1:5000 dilution.  
Lysates/proteins: 50ng - 100ng per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.



Western blot analysis of lysates from Rat IgG, using HRP Goat Anti-Rat IgG (H+L) (AS028) at 1:1000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.