

FITC-conjugated Donkey anti-Rabbit IgG (H+L)

Catalog No.: AS042 6 Publications

Basic Information

Observed MW

Calculated MW

Category

Secondary antibody

Applications

IF/ICC,FC

Cross-Reactivity

Rabbit

Conjugate

FITC. Ex:491nm. Em:516nm.

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

IF/ICC 1:50 - 1:200

FC 1:50 - 1:200

Immunogen Information

Gene ID Swiss Prot

Immunogen

This information is considered to be commercially sensitive.

Synonyms

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn

Product Information

SourceIsotypePurificationDonkeyFluorescein conjugated IgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.025% Sodium Azide, 0.75% BSA, 50% glycerol, pH7.3.

Validation Data






Flow cytometric analysis of Positive antibody E-Cadherin Rabbit mAb (2.5µg/mL) in various cells (orange) compare to Rabbit rabbit isotype control (blue) and non-staining control (Red). The secondary antibody used was FITC Donkey Anti-Rabbit IgG (H+L) (AS042) at 1:100.



Immunofluorescence analysis of PC-12 cells using Vimentin Rabbit mAb (A19607) at a dilution of 1:100 (40x lens). Secondary antibody: FITC Donkey Anti-Rabbit IgG (H+L) (AS042) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using Vimentin Rabbit mAb (A19607) at a dilution of 1:100 (40x lens). Secondary antibody: FITC Donkey Anti-Rabbit IgG (H+L) (AS042) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.