

ABflo® 647-conjugated F(ab')₂ Fragment Goat anti-Rabbit IgG, Fc fragment specific

Catalog No.: AS086 2 Publications

Basic Information

Observed MW

Calculated MW

Category

Secondary antibody

Applications

IF-P,FC

Cross-Reactivity

Rabbit

Conjugate

ABflo® 647, Ex:648nm, Em:664nm,

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

IF-P 1:100 - 1:500

FC 1:100 - 1:800

Immunogen Information

Gene ID Swiss Prot

Immunogen

This information is considered to be commercially sensitive.

Synonyms

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn

Product Information

SourceIsotypePurificationGoatIgGAffinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.025% Sodium Azide, 0.75% BSA, 50% glycerol, pH7.3.

Validation Data







Flow cytometry: Daudi cells(+) and Jurkat cells(-) were stained with Rabbit IgG isotype control (AC042, 10 μ g/mL, blue line) or CD20 Rabbit mAb (A4893, 10 μ g/mL orange line), followed by Goat anti-Rabbit pAb ABflo® 647 (AS086, 1:600 dilution) staining. Nonfluorescently stained Daudi cells was used as blank control (red line).

Confocal imaging of paraffin-embedded Mouse brain using β III-Tubulin Rabbit mAb (A17913, dilution 1:200) followed by a further incubation with ABflo® 647 F(ab')² Fragment Goat Anti-Rabbit IgG, Fc fragment specific(AS086, dilution 1:500)(Red). DAPI was used for nuclear staining (Blue). Objective: 40x.Perform high pressure antigen retrieval with 0.01M citrate buffer (pH 6.0) prior to IF staining.