# ABclonal www.abclonal.com

# ABflo® 647 Rabbit anti Mouse IgG3 (Fc) mAb

Catalog No.: AS152

### **Basic Information**

# **Background**

**Observed MW** 

**Calculated MW** 

36kDa

Category

Secondary antibody

**Applications** 

FC

**Cross-Reactivity** 

Mouse

CloneNo number

ARC68336

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

## **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

# **Immunogen Information**

Gene ID

Swiss Prot P03987-2

**Immunogen** 

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

**Synonyms** 

#### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Т	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification


**Storage** 

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

### **Validation Data**





Flow cytometry: 1X10^6 THP-1 cells (negative control,left) and BALB/c mouse splenocytes (right) were surface-stained with Purified anti-mouse I-Ad Antibody (2  $\mu g/mL$ ,orange line) or Mouse IgG isotype control (2  $\mu g/mL$ ,blue line), followed by ABflo® 647 Rabbit anti Mouse IgG3 (Fc) mAb (AS152,5  $\mu l/Test$ ) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 BALB/c mouse splenocytes were surface-stained with Mouse IgG isotype control (2 µg/mL,Ieft) or Purified anti-mouse I-Ad Antibody (2 µg/mL,right). ABflo® 647 Rabbit anti Mouse IgG3 (Fc) mAb (AS152,5 µl/Test) was used as a secondary antibody.