

# PE Rabbit anti-Mouse IgG (Fc) mAb

Catalog No.: AS154

## Basic Information

### Observed MW

### Calculated MW

36kDa

### Category

Secondary antibody

### Applications

FC

### Cross-Reactivity

Mouse

### CloneNo number

ARC68493

### Conjugate

PE. Ex:565nm. Em:574nm.

## Background

## Recommended Dilutions

**FC** 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Immunogen Information

### Gene ID

/

### Swiss Prot

P01868

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  Raji cells (negative control, left) and Human PBMC (right) were surface-stained with Mouse anti Human CD3 mAb (2  $\mu\text{g}/\text{mL}$ , orange line) or Mouse IgG isotype control (2  $\mu\text{g}/\text{mL}$ , blue line), followed by PE Rabbit anti-Mouse IgG (Fc) mAb (AS154, 5  $\mu\text{l}/\text{Test}$ ) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  Human PBMC were surface-stained with Mouse IgG isotype control (2  $\mu\text{g}/\text{mL}$ , left) or Mouse anti Human CD3 mAb (2  $\mu\text{g}/\text{mL}$ , right). PE Rabbit anti-Mouse IgG (Fc) mAb (AS154, 5  $\mu\text{l}/\text{Test}$ ) was used as a secondary antibody.