# ABclonal® www.abclonal.com

# CDH2 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM01772

## **Basic Information**

#### Catalog No.

RM01772

#### Category

Cell Lysate

# **Parental Cell line**

HeLa

#### Genotype

Knockout

# **Background**

This gene encodes a classical cadherin and member of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein is proteolytically processed to generate a calcium-dependent cell adhesion molecule and glycoprotein. This protein plays a role in the establishment of left-right asymmetry, development of the nervous system and the formation of cartilage and bone. [provided by RefSeq, Nov 2015]

## **Gene Information**

#### **Species**

Human

# Gene ID

1000

#### **Swiss Prot**

P19022

#### **Synonyms**

CD325; CDHN; CDw325; NCAD

# **Contact**

8	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Product Information**

#### **Description**

CDH2 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:44bp deletion in exon2

Allele-2:44bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

# **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

# **Shipping Conditions**

**Amount** 

4°C

50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

# Sequencing data

WT AAAGAGACCCAGGA\*\*\*\*\*\*\*\*\*\*\*\*\*CTGAGGAGTCAGTG
Mut AAAGAGACCCAGGA\*\*\*Deletion\*\*\*CTGAGGAGTCAGTG
Allele-1: 44bp deletion in exon2

WT AAAGAGACCCAGGA\*CTGAGGAGTCAGTG
Mut AAAGAGACCCAGGA\*\*\*Deletion\*\*\*\*CTGAGGAGTCAGTG
Allele-2: 44bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CDH2 knockout (KO) HeLa cells, using sanger sequencing.