

CTNNB1 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM01773

Basic Information

Catalog No.

RM01773

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

CTNNB1

Species

Human

Gene ID

1499

Swiss Prot

P35222

Synonyms

CTNNB; MRD19; armadillo

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2016]

Product Information

Description

CTNNB1 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:43bp deletion in exon2

Allele-2:5bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CATGGCCATGGAAC*****AGAAAAGCGGCTG
Mut CATGGCCATGGAAC***Deletion***AGAAAAGCGGCTG
Allele-1: 5 bp deletion in exon2

WT ATGGAACCAGACA*****CTCTGGAATCCAT
Mut ATGGAACCAGACA***Deletion***CTCTGGAATCCAT
Allele-2: 43 bp deletion in exon2

WT CCATGGAACCAG*****CAGTCCT
Mut CCATGGAACCAG**Insertion*Deletion***CAGTCCT
Allele-3: 63bp Insertion and 164 bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CTNNB1 knockout (KO) 293T cells, using sanger sequencing.