

# EGLN1 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM01800 **1 Publications**

## Basic Information

**Catalog No.**

RM01800

**Category**

Cell Lysate

**Parental Cell line**

293T

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

EGLN1

**Species**

Human

**Gene ID**

54583

**Swiss Prot**

Q9GZT9

**Synonyms**C1orf12; ECTY3; HALAH; HIF-PH2;  
HIFPH2; HPH-2; HPH2; PHD2; SM20;  
ZMYND6

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The protein encoded by this gene catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. HIF is a transcriptional complex that plays a central role in mammalian oxygen homeostasis. This protein functions as a cellular oxygen sensor, and under normal oxygen concentration, modification by prolyl hydroxylation is a key regulatory event that targets HIF subunits for proteasomal destruction via the von Hippel-Lindau ubiquitylation complex. Mutations in this gene are associated with erythrocytosis familial type 3 (ECYT3).

## Product Information

**Description**

EGLN1 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:73bp deletion in exon1

Allele-2:74bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**

4°C

**Amount**

50μL, 2μg/μL.

**Storage**

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

**Protocol**


To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

WT   GCCGGGACAACGCC\*\*\*\*\*GTGCGGCCGCCGGC  
Mut   GCCGGGACAACGCC\*\*\*Deletion\*\*\*GTGCGGCCGCCGGC  
Allele-1: 73bp deletion in exon1  
  
WT   CGCCGGGACAACGCC\*\*\*\*\*GTGCGGCCGCCGGC  
Mut   CGCCGGGACAACGCC\*\*\*Deletion\*\*\*GTGCGGCCGCCGGC  
Allele-2: 74bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and EGLN1 knockout (KO) 293T cells, using sanger sequencing.

## WB data



Western blot analysis of extracts from parental (Control) and EGLN1 knockout (KO) 293T cells, using EGLN1 antibody (A14557) at 1:3000 dilution.