

SIRT3 Knockout 293T Cell Line, Homozygous

Catalog No.: RM01802

Basic Information

Catalog No.

RM01802

Category

Cell Line

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

SIRT3

Species

Human

Gene ID

23410

Swiss Prot

Q9NTG7

Synonyms

SIR2L3

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Two alternatively spliced transcript variants that encode different proteins have been described for this gene.

Product Information

Description

SIRT3 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:128bp deletion in exon2

Allele-2:128bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial.

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.


1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT CGATCTCCCGTACC*****AAGGGGCTGCTTCT
Mut CGATCTCCCGTACC***Deletion***AAGGGGCTGCTTCT
Allele-1: 128bp deletion in exon2
WT CGATCTCCCGTACC*****AAGGGGCTGCTTCT
Mut CGATCTCCCGTACC***Deletion***AAGGGGCTGCTTCT
Allele-2: 128bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and SIRT3 knockout (KO) 293T cells, using sanger sequencing.

WB data



Western blot analysis of extracts from parental (Control) and SIRT3 knockout (KO) 293T cells, using SIRT3 antibody (A5718) at 1:1000 dilution.